

University of Groningen

## Emissiespectra van lichte elementen en hun verbindingen in het ultra-zachte Röntgegebied

Hof, Sietze

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*

Publisher's PDF, also known as Version of record

*Publication date:*

1940

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Hof, S. (1940). Emissiespectra van lichte elementen en hun verbindingen in het ultra-zachte Röntgegebied. Groningen: Noordhoff Uitgevers.

**Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

**Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

## SUMMARY.

The K-emission spectra of Be, B and C, and the L-emission spectra of Al and Si of the pure elements in the solid state and some compounds are investigated. The spectra exist of broad lines, due to the broad energy-level, which is occupied by the valency-electrons. As to the pure elements, it could be understood, why the insulators have emission lines which gradually diminish in intensity at both sides (fig. 18). The conductors, however, have emission lines which at the short wave-length side have a sharp edge (the intensity falls suddenly from a given value to zero). This edge corresponds with the energy up to which the valency-band is filled (fig. 5, fig. 8, fig. 13, fig. 16). It is of interest to notice that it may be concluded from the K-line of B that this element has to be a conductor.

It is pointed out that in the case of a compound not only the density of the energy states in the valency-band but also the transition probability plays a role in determining the form of the emission line. In this way we can understand why the beryllium K-lines of BeO (fig. 6) and BeSO<sub>4</sub> (fig. 7) are nearly identical, whereas the crystal lattices of these compounds are totally different. The same can be said of B<sub>2</sub>O<sub>3</sub> (fig. 10) and Na<sub>2</sub>B<sub>4</sub>O<sub>7</sub> (fig. 11).

Some special questions are discussed in detail. The carbon K-line in SiC has a sharp edge, this is in agreement with the fact that SiC is a conductor.

A new diffuse and faint emission line of Be in BeO lying on the long wave-length side of the ordinary K-line of this element has been found. It is interpreted as a  $K \rightarrow L$  transition.